

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 9/20, 31/565, 47/00	A1	(11) International Publication Number: WO 95/06461 (43) International Publication Date: 9 March 1995 (09.03.95)
(21) International Application Number: PCT/US94/09945 (22) International Filing Date: 2 September 1994 (02.09.94) (30) Priority Data: 9318348.1 3 September 1993 (03.09.93) GB 9323864.0 19 November 1993 (19.11.93) GB 9411903.9 14 June 1994 (14.06.94) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; Corporate Intellectual, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FAULKNER, Patrick, Gerard [IE/US]; 7008 Shawnee Circle, Collegeville, PA 19426 (US). FISHER, Mark, Warren [AU/US]; 484 Keebler Road, King of Prussia, PA 19406 (US). SAUER, Joseph, Peter [US/US]; 130 Ohio Avenue, Clementin, NJ 08021 (US). HERNANDEZ, Carlos, Roberto [US/US]; 7481 Grant Avenue, Pennsauken, NJ 08109-3151 (US).		(74) Agents: DUSTMAN, Wayne, J. et al; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US). (81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: STABILIZED TABLET FORMULATION (57) Abstract Invented are stabilized tablet formulations of substituted 3,5-diene steroidal compounds. Also invented are methods of preparing stabilized tablet formulations of substituted 3,5-diene steroidal compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MD	Republic of Moldova	TJ	Tajikistan
DE	Germany	MG	Madagascar	TT	Trinidad and Tobago
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain	MN	Mongolia	US	United States of America
FI	Finland			UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				


STABILIZED TABLET FORMULATION

The present invention relates to novel stabilized tablet formulations of substituted 3,5-diene steroidal compounds containing a 3 position electron-withdrawing group and methods of preparing stabilized tablet formulations of substituted 3,5-diene steroidal compounds containing a 3 position electron-withdrawing group. Preferably, this invention relates to novel stabilized tablet formulations of substituted 3,5-diene steroidal compounds containing a 3 position acid group and methods of preparing stabilized tablet formulations of substituted 3,5-diene steroidal compounds containing a 3 position acid group. Preferably, this invention relates to novel stabilized tablet formulations of substituted 3,5-diene steroidal compounds containing a 3 position carboxylic acid group and methods of preparing stabilized tablet formulations of substituted 3,5-diene steroidal compounds containing a 3 position carboxylic acid group. Advantageously, the substituted 3,5-diene steroidal compound for use in the presently invented stabilized tablet formulations and presently invented methods is N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid.

Detailed Description of the Invention

Novel substituted 3,5-diene steroidal compounds containing a 3 position electron-withdrawing group have recently been shown to exhibit significant pharmaceutical activity. Specifically, said compounds have been found to inhibit steroid 5- α -reductase and are thereby useful in the treatment of prostate diseases such as benign prostatic hypertrophy. Stabilized tablet formulations of substituted 3,5-diene steroidal compounds containing a 3 position electron-withdrawing group are advantageous in preparing dosage units for oral administration.

By the term "substituted 3,5-diene steroidal compound" as used herein is meant a compound of the figure



wherein W is an electron-withdrawing group; and the C₁ and C₂ positions and the C₈ to C₁₉ positions are optionally substituted with pharmaceutically acceptable

substituents, degrees of unsaturation or a combination of pharmaceutically acceptable substituents and degrees of unsaturation.


5 The term "electron-withdrawing group" is well known to those of skill in the art and is used herein as its standard meaning which is a functional group that draws electrons to itself more than a hydrogen atom would if it occupied the same position in the molecule, as described in J. March, *Advanced Organic Chemistry*, third edition, Pub: John Wiley & Sons, Inc. (1985). Preferred electron-
10 withdrawing groups for use in the presently invented stabilized tablet formulations and presently invented methods are: $-\text{COOH}$, $-\text{P}(\text{O})(\text{OH})_2$, $-\text{PH}(\text{O})\text{OH}$, $-\text{SO}_3\text{H}$, $-\text{CHO}$, $-\text{CONHSO}_2\text{R}^{20}$ where R^{20} is an alkyl group having from 1 to 6 carbon atoms, $-\text{CN}$, $-\text{CONH}_2$ and, where appropriate, esters thereof.

15 By the term "acid group" as used herein is meant a substituent selected from the group consisting of: $-\text{COOH}$, $-\text{P}(\text{O})(\text{OH})_2$, $-\text{PH}(\text{O})\text{OH}$ and $-\text{SO}_3\text{H}$.

The most preferred electron-withdrawing group for use in the presently invented stabilized tablet formulations and presently invented methods is $-\text{COOH}$.

20 By the term "fatty acid" as used herein is meant saturated and unsaturated fatty acids of the standard usage wherein the compound contains from about 10 to 22 carbon atoms, preferably decanoic acid, stearic acid, palmitic acid, oleic acid, lauric acid or myristic acid; most preferably, stearic acid.

25 Preferred substituted 3,5-diene steroidal compounds for use in the presently invented stabilized tablet formulations and presently invented methods include compounds of the Formula I




wherein R is a hydrocarbon radical selected from substituted or unsubstituted 1- or 2-adamantyl, 1-, 2- or 7-norbornanyl.

- Included within the scope of compounds of Formula I are those wherein the
- 5 1- or 2-adamantyl or 1-, 2- or 7-norbornanyl moieties are substituted with one or more of: C₁-C₄ linear or branched alkyl; nitro; oxo; C₇-C₉ arylalkyl, including benzyl; (CH₂)_nCOOR¹ where n is 0-2 and R¹ is H or linear or branched C₁-C₄ alkyl; CH₂OH; OH; OR² where R² is C₁-C₄ linear or branched alkyl; halo; CONH₂; CH₂NH₂; CH₂NHCOR³ where R³ is C₁-C₄ linear or branched alkyl;
- 10 phenyl and p-substituted phenyl wherein the substituents are members selected from the group consisting of nitro; amino; sulfo and cyano.


- Compounds of Formula I and methods of preparing compounds of Formula I are described in EPO Publn. No. 0465141 (Published 8 January 1992) to
- 15 Rasmusson, et al. (Merck Co. Inc.). Said compounds are disclosed therein as being useful in inhibiting steroid 5- α -reductase.

- Preferred substituted 3,5-diene steroidal compounds for use in the presently invented stabilized tablet formulations and presently invented methods include
- 20 compounds of the Formula II



- wherein R is C₁-C₆ linear or branched alkyl; C₃-C₈ cycloalkyl, which can be substituted with C₁-C₄ alkoxy or C₁-C₄ linear or branched alkyl; C₆-C₁₂ aryl, which can be substituted with one or more of: -OH, -OC₁-C₄ alkyl, C₁-C₄ alkyl,
- 25 -(CH₂)_mOH, -(CH₂)_nCOOH, including protected -OH, where m is 1-4 and n is 1-3.


Additionally preferred substituted 3,5-diene steroidal compounds for use herein are the ketone reduction products of II, the secondary alcohols of the formula



wherein R is C₁-C₆ linear or branched alkyl; C₃-C₈ cycloalkyl, which can be substituted with C₁-C₄ alkoxy or C₁-C₄ linear or branched alkyl; C₆-C₁₂ aryl, which can be substituted with one or more of: -OH, -OC₁-C₄ alkyl, C₁-C₄ alkyl, -(CH₂)_mOH, -(CH₂)_nCOOH, including protected -OH, where m is 1-4 and n is 1-3.

Compounds of Formula II and the ketone reduction products of II and methods of preparing compounds of Formula II and the ketone reduction products thereof are described in EPO Publ. No. 0465142 (published 8 January 1992) to Rasmusson, et al. (Merck & Co. Inc.). Said compounds and ketone reduction products thereof are disclosed therein as being useful in inhibiting steroid 5- α -reductase.

Particularly preferred substituted 3,5-diene steroidal compounds for use in the presently invented stabilized tablet formulations and presently invented methods include compounds of the Formula III



in which;

R² is H or CH₃ and

R³ is

(1) α -hydrogen, α -hydroxyl, or α -acetoxy and/or

(a)



where W is a bond or C₁₋₁₂alkyl, and R⁴ is

- (i) hydrogen,
 (ii) hydroxyl,
 (iii) C₁₋₈alkyl,
 5 (iv) hydroxy C₁₋₈alkyl,
 (v) C₁₋₈alkoxy,
 (vi) NR⁵R⁶, where R⁵ and R⁶ are each independently selected from hydrogen, C₁₋₈alkyl, C₃₋₆cycloalkyl, phenyl; or R⁵ and R⁶ taken together with the nitrogen to which they are attached represent
 10 a 5-6 membered saturated ring comprising up to one other heteroatom selected from oxygen and nitrogen, or
 (vii) OR⁷, where R⁷ is hydrogen, alkali metal, C₁₋₁₈alkyl, benzyl, or
- (b) β-Alk-OR⁸, where Alk is C₁₋₁₂alkyl, and R⁸ is
 (i) phenylC₁₋₆alkylcarbonyl,
 (ii) C₅₋₁₀cycloalkylcarbonyl,
 (iii) benzoyl,
 (iv) C₁₋₈alkoxycarbonyl,
 20 (v) aminocarbonyl, or C₁₋₈alkyl substituted aminocarbonyl,
 (vi) hydrogen, or
 (vii) C₁₋₈alkyl,

- (2) =CH-W-CO-R⁴ or =CH-W-OR⁸, where W is a bond or C₁₋₁₂alkyl and R⁴ and R⁸ have the same meaning as above and R⁸ also is hydrogen or C₁₋₂₀alkylcarbonyl;

(3)



- 30 where the dashed bond replaces the 17-α -hydrogen,

- (4) α -hydrogen and β-NHCOR⁹ where R⁹ is C₁₋₁₂alkyl or β-NR⁵R⁶ where R⁵ and R⁶ have the same meaning as above,


- 35 (5) α -hydrogen and β-cyan ,

(6) α -hydrogen and β -tetrazolyl, r

(7) keto.

5 Compounds of Formula III and methods of preparing compounds of Formula III are described in U.S. Patent No. 5,017,568. Said compounds are disclosed therein as being useful in inhibiting steroid 5- α -reductase.






10 Particularly preferred substituted 3,5-diene steroidal compounds for use in the presently invented stabilized tablet formulations and presently invented methods include compounds of the formula (IV)






wherein

Y is oxygen or sulphur,

15 R_1 is methyl, ethyl, isopropyl, tert-butyl, cyclohexyl; and the group




20



; or

Compounds of formula IV and methods of preparing compounds of formula IV are described in U.S. Patent No. 5,212,166. Said compounds are disclosed therein as being useful in inhibiting steroid 5 α -reductase.

- 5 Particularly preferred substituted 3,5-diene steroidal compounds for use in the presently invented stabilized tablet formulations and presently invented methods include compounds of the formula (V)



wherein:

- 10 R^1 represents: a hydrogen atom; an alkyl group having from 1 to 6 carbon atoms; or a substituted alkyl group having from 1 to 6 carbon atoms and having at least one substituent selected from aryl groups as defined below and aromatic heterocyclic groups as defined below;
- R^2 represents:
- 15 a substituted alkyl group having from 1 to 6 carbon atoms and having at least one substituent selected from aryl groups as defined below and aromatic heterocyclic groups as defined below, and said alkyl group further optionally having a single hydroxy or carboxy substituent; or
- a diarylamino group in which the two aryl parts are the same or different and
- 20 each is as defined below;
- R^3 represents a carboxy group or a group of formula $-\text{CONHSO}_2\text{R}^4$ wherein R^4 represents an alkyl group having from 1 to 6 carbon atoms;
- said aryl groups are carbocyclic aromatic groups having from 6 to 14 ring carbon atoms and which are unsubstituted or are substituted by at least one substituent
- 25 selected from substituents A, defined below;
- said aromatic heterocyclic groups have 5 or 6 ring atoms of which from 1 to 3 are hetero-atoms selected from nitrogen, oxygen and sulphur hetero-atoms and the remainder are carbon atoms, said group being unsubstituted or being substituted by at least one substituent selected from substituents B, defined below;
- 30 said substituents A are selected from: alkyl groups having from 1 to 6 carbon atoms; alkoxy groups having from 1 to 6 carbon atoms; alkoxycarbonyl groups having from 2 to 7 carbon atoms; hydroxy groups; halogen atoms; amino groups; alkylamino

groups having from 1 to 6 carbon atoms; dialkylamino groups in which each alkyl part has from 1 to 6 carbon atoms; aliphatic acylamino groups having from 1 to 6 carbon atoms; aromatic acylamino groups in which the aromatic part is a carbocyclic aryl group which has from 6 to 10 ring carbon atoms and which is unsubstituted or is substituted by at least one substituent selected from substituents C, defined below; cyano groups; nitro groups; and carboxy groups; said substituents B are selected from: alkyl groups having from 1 to 6 carbon atoms; alkoxy groups having from 1 to 6 carbon atoms; hydroxy groups; halogen atoms; carbocyclic aryl groups which have from 6 to 10 ring carbon atoms and which are unsubstituted or are substituted by at least one substituent selected from substituents C, defined below: amino groups; alkylamino groups having from 1 to 6 carbon atoms; dialkylamino groups in which each alkyl part has from 1 to 6 carbon atoms; aliphatic acylamino groups having from 1 to 6 carbon atoms; aromatic acylamino groups in which the aromatic part is a carbocyclic aryl group which has from 6 to 10 ring carbon atoms and which is unsubstituted or is substituted by at least one substituent selected from substituents C, defined below; nitro groups; and carboxy groups; said substituents C are selected from: alkyl groups having from 1 to 6 carbon atoms; alkoxy groups having from 1 to 6 carbon atoms; hydroxy groups; halogen atoms; amino groups; alkylamino groups having from 1 to 6 carbon atoms; dialkylamino groups in which each alkyl part has from 1 to 6 carbon atoms; aliphatic acylamino groups having from 1 to 6 carbon atoms; cyano groups; nitro groups; and carboxy groups.

Compounds of formula V and methods of preparing compounds of formula V are described in European Patent Application Publication Number 0567271 A2, published October 27, 1993. Said compounds are disclosed therein as being useful in inhibiting steroid 5 α -reductase.

As used herein, unless otherwise specified, C_{1-n} alkyl and C_{1-n} alk means a straight or branched hydrocarbon chain having 1 to n carbons and alk means a straight or branched hydrocarbon chain having 1 to 12 carbons.

Preferably the substituted 3,5-diene steroidal compound for use in the presently invented stabilized tablet formations and presently invented methods is N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid. N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid can be prepared as described in U.S.

Patent No. 5,017,568. This compound is disclosed therein as being useful in inhibiting steroid 5 α reductase.

Although stable when blended or granulated with one or more commonly used pharmaceutical excipients such as, for example, lactose, tribasic calcium phosphate, silicon dioxide, corn starch, sodium starch glycolate, hydroxypropyl cellulose, hydroxypropyl methylcellulose, gelatin, magnesium stearate, microcrystalline cellulose, mannitol, and sorbitol, substituted 3,5-diene steroidal compounds were unexpectedly found to undergo a significant increase in the rate of formation of degradation products when compressed into tablets with such pharmaceutical excipients. This increase in degradation products occurs whether the substituted 3,5-diene steroidal compound and said excipients are formulated in direct compression tablets or in wet granulation tablets.

As used herein, unless otherwise indicated, the term "tablet" and derivatives thereof, means a direct compression tablet or a wet granulation tablet.

The reduction of degradation products of steroidal compounds in pharmaceutical formulations is of particular importance because minor changes in the structure of steroids are known to produce profound changes in its biological activity. Moreover, steroidal compounds are known to be extremely potent, often requiring only very low doses of the compound to exhibit pharmaceutical activity.

It has now been found that the addition of a fatty acid, preferably stearic acid or palmitic acid; an antioxidant, preferably butylated hydroxyanisole (BHA) or butylated hydroxytoluene (BHT), most preferably BHT; or a combination of a fatty acid and an antioxidant, to a blend of pharmaceutical excipients containing a substituted 3,5-diene steroidal compound dramatically decreases the percentage of degradation products of the active in subsequently compressed tablets. Said tablets are prepared by compression under normal operating conditions using a standard tablet press.

The stability of N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid (compound A) in various tablet formulations was determined by HPLC analysis. Degradation products were analyzed by LC/MS and unexpectedly found to be the result of oxidative additions (the 7 keto oxidation product and others) which only occurred when blends or granulations of Compound A were

compressed into tablets. In order to reduce the percentage of degradation products to desired levels, preferably an increase of no more than 0.2 percent from baseline at day 0 after two weeks in the Accelerated Stability Screening Method (described herein), numerous tablet formulations were prepared and analyzed as indicated
5 below.

In order to facilitate research efforts an Accelerated Stability Screening Method was developed. It was found that storing tablets at 85°C for a short time (preferably two weeks) is suitable for predicting long term stability at room
10 temperature. This method was employed upon noting that the degradation profiles generated at room temperature as compared to 85°C were similar and that the profiles generated for tablets prepared by either direct compression (dry granulation process) or fluid bed granulation (wet granulation process) are the same.

15 Direct Compression Tablets

Experiments 1 and 2 were performed (utilizing Compound A) in order to determine the most advantageous direct compression tablet formulations of substituted 3,5-diene steroidal compounds.

20 Experiment 1

In Experiment 1 Compound A, microcrystalline cellulose, lactose, corn starch and sodium starch glycolate shown in Table I below, were mixed in the proportions shown. The mixture was split in half (into Batch 1 and Batch 2). Batch 1 was mixed with 1 percent magnesium stearate and compressed into
25 appropriately sized tablets preferably 300mg to 350mg tablets each containing from about 80mg to 85mg of active compound (no stearic acid group). Batch 2 was first mixed and granulated with 2 percent stearic acid, then mixed and granulated with 1 percent magnesium stearate and compressed into appropriately sized tablets preferably 300mg to 350mg tablets each containing from about 80mg to 85mg of
30 active compound (stearic acid group). The tablets thus prepared were placed in stability chamber at 85° and the results of the stability test were recorded in Table 2 below.

Table 1

		%w/w
Batch 1	Compound A	10
	Microcrystalline Cellulose	40
	Lactose	32
	Corn Starch	10
	Sodium Starch Glycolate	7
	Mg Stearate	1
Batch 2	Compound A	10
	Microcrystalline Cellulose	40
	Lactose	32
	Corn Starch	10
	Sodium Starch Glycolate	7
	Stearic Acid	+2
	Mg Stearate	1

Table 2

	% Degradation	% Degradation	% Degradation
	Day 0	Day 4	Day 11
Batch 1 (No Stearic Acid)	0	0.05	0.17
Batch 2 (Stearic Acid)	0	0.01	0.01

5

Experiment 2

In Experiment 2 one gram of Compound A was mixed and granulated with ten grams of lactose. The mixture was split into three equal parts (Batch 3 Batch 4 and Batch 5). Batch 3 was compressed into tablets of about 300mg to 350mg, about 30mg to 35mg of active compound (Control group: no additives). Batch 4 was mixed with two percent pure stearic acid and compressed into tablets of about 300mg to 350mg, about 30mg to 35mg of active compound. Batch 5 was mixed with two percent pure palmitic acid and compressed into tablets of about 300mg to

10

350mg, about 30mg to 35mg of active compound. The tablets thus prepared were placed in stability chamber at 85°C and the results of the stability test were recorded in Table 3 below.

5

Table 3

	% Degradation Day 0	% Degradation Day 6	% Degradation Day 11
Batch 3 (Control)	0	0.47	0.54
Batch 4 (Pure Stearic Acid)	0	0.03	0.04
Batch 5 (Pure Palmitic Acid)	0	0.08	0.11

Wet Granulation Process

Experiments 3 to 6 were performed (utilizing Compound A) in order to determine the most advantageous wet granulation tablet formulations of substituted 3,5-diene compounds. To perform the wet granulation experiments a 'Standard Granulation' containing Compound A was utilized. To prepare the Standard Granulation Compound A (generally about 37% of this mixture) and lactose (generally about 57% of this mixture) are mixed and granulated with about a 9% solution of hydroxypropylmethyl cellulose (generally about 6% of this mixture). The wet granules are screened and dried to form the Standard Granulation.

10

15

Experiment 3

In Experiment 3 Standard Granulation, corn starch, sodium starch glycolate, microcrystalline cellulose, magnesium stearate and stearic acid (for Batch 6) or additional magnesium stearate (for Batch 7) shown in Table 4 below, were blended in the proportions shown and compressed into tablets of about 300mg to 350mg, about 80mg to 85mg of active compound. The tablets thus prepared were placed in stability chamber at 85°C and the results of the stability test were recorded in Table 5 below.

20

25

Table 4

Batch 6	
Standard Granulation	715 g
Stearic Acid	23 g
Corn Starch	49.5 g
Sodium Starch Glycolate	50 g
Microcrystalline Cellulose	150 g
Magnesium Stearate	12.5 g
Batch 7	
Standard Granulation	715 g
Magnesium Stearate	23 g
Corn Starch	49.5 g
Sodium Starch Glycolate	50 g
Microcrystalline Cellulose	150 g
Magnesium Stearate	12.5 g

Table 5

	% Degradation Day 0	% Degradation Day 11
Batch 6	0	0.13
Batch 7	0	0.31

5

Experiment 4

In Experiment 4 Standard Granulation and BHT (for Batch 8) shown in Table 6 below, or stearic acid and BHT (for Batch 9) shown in Table 7 below, were mixed and granulated in the proportions shown and compressed into tablets of about 350 mg total tablet weight. The tablets thus prepared were placed in stability chamber at 85°C and the results of the stability test were recorded in Tables 6 and 7 below.

10

Table 6

Batch 8		
% w/w BHT	% Degradation	% Degradation
	Day 0	Day 14
0	0	0.57
0.1	0	0.22
0.5	0	0.45
1.0	0	0.17
2.0	0	0.10

Table 7

Batch 9			
%w/w Stearic Acid	% w/w BHT	% Degradation	% Degradation
		Day 0	Day 14
0.2	0.1	0	0.34
1.0	0.5	0	0.04
2.0	1.0	0	0.02
4.0	2.0	0	0.03

5

Experiment 5

In Experiment 5 Standard Granulation (at about 72% w/w), microcrystalline cellulose (from 14% to 17% w/w depending on BHT and stearic acid content from Table 8) corn starch (at about 5% w/w), sodium starch glycolate (at about 5% w/w) and stearic acid and BHT (for Batch 10) shown in Table 8 below, were mixed and granulated in the proportions shown and compressed into 300 mg tablets (about 80 mg of Compound A). The tablets thus prepared were placed in stability chamber at 85°C and the results of the stability test were recorded in Table 8 below.

10

Table 8

Batch 10			
%w/w Stearic Acid	% w/w BHT	% Degradation	% Degradation
		Day 0	Day 15
0	0	0	0.75
0	0	0	0.59
0.5	0.25	0	0.1
1.0	0.52	0	0
1.6	0.79	0	0.03
2.0	1.05	0	0.01
2.0	1.05	0	0.03

Experiment 6

- 5 In Experiment 6 Standard Granulation (at about 72% w/w), microcrystalline cellulose (from 16% to 17% w/w depending on BHT content from Table 9), corn starch (at about 5% w/w), sodium starch glycolate (at about 5%) magnesium stearate (at about 1%), stearic acid (at about 2%) and BHT (for Batch 11) shown in Table 9 below, were mixed and granulated in the proportions shown and compressed into 300 mg tablets (about 80 mg of Compound A). The tablets thus prepared were placed in stability chamber at 85°C and the results of the stability test were recorded in Table 9 below.
- 10

Table 9

Batch 11			
%w/w BHT	% Degradation	% Degradation	% Degradation
	Day 0	Day 7	Day 16
0	0	0.12	0.19
0.1	0	0	0
0.25	0	0	0
0.5	0	0	0
1.0	0	0.01	0.03

- 15 The data from the above experiments demonstrates the ability of the presently discovered invention to stabilize direct compression tablets and wet granulation tablets containing substituted 3,5-diene steroidal compounds.

Fatty acids, particularly stearic acid, are known pharmaceutical excipients which are utilized in the art, primarily as tablet lubricants. Stearic acid, as a tablet lubricant, is considered interchangeable with other known tablet lubricants, particularly magnesium stearate. As discovered herein, fatty acids are disclosed as exhibiting the previously unknown ability to stabilize tablet formulations of substituted 3,5-diene steroidal compounds when other known tablet lubricants, particularly magnesium stearate, and other known pharmaceutical excipients do not exhibit such activity. As such, the present invention also relates to the use of fatty acids, preferably stearic acid or palmitic acid, most preferably stearic acid, as stabilizing agents in tablet formulations, particularly tablet formulations of substituted 3,5-diene steroidal compounds.

Preferably the fatty acid will be present in the finished tablet, as prepared herein, in an amount from about 0.5% to about 10% by weight.

Most preferably the fatty acid will be present in the finished tablet, as prepared herein, in an amount from about 1% to about 3% by weight, most preferably about 2%.

By the term "stabilized", and derivatives thereof, as used herein is meant that the subject substituted 3,5-diene steroidal compound when compressed into a tablet, under normal operating conditions using a standard tablet press, with a blend of pharmaceutical excipients and a fatty acid, an antioxidant or a combination of a fatty acid and an antioxidant will experience the formation of less degradation products, preferably less than one half of the degradation products, experienced by said substituted 3,5-diene steroidal compound when compressed into a tablet with the same blend of pharmaceutical excipients without such fatty acid, such antioxidant or such combination of a fatty acid and an antioxidant. Assessing the ability of a fatty acid, an antioxidant or a combination of a fatty acid and an antioxidant to stabilize a particular substituted 3,5-diene steroidal compound is accomplished by comparing the stability of tablets containing a blend of pharmaceutical excipients, the subject substituted 3,5-diene steroidal compound and such fatty acid, such antioxidant or such combination of a fatty acid and an antioxidant to tablets wherein the only difference is that said fatty acid, said antioxidant or said combination of a fatty acid and an antioxidant is left out (as in Experiment 1) or substituted for by another pharmaceutical excipient, such as magnesium stearate (as in Experiment 3).

This invention provides for a method of stabilizing tablet formulations of substituted 3,5-diene steroidal compounds which comprises:


- (a) admixing a fatty acid, an antioxidant or a combination of a fatty acid and an antioxidant with a pharmaceutical excipient blend and a substituted 3,5-diene steroidal compound;
- (b) compressing the resulting mixture into tablets.

This invention also provides for a pharmaceutical formulation in tablet form comprising: a pharmaceutical excipient blend, a substituted 3,5-diene steroidal compound and a fatty acid, an antioxidant or a combination of a fatty acid and an antioxidant, preferably stearic acid for a direct compression tablet; preferably a combination of stearic acid and BHT for a wet granulation tablet.

By the term "pharmaceutical excipient blend" and derivatives thereof, as used herein is meant a mixture, a granulation or an admix of a mixture and a granulation of a plurality of commonly used pharmaceutical excipients, such as lactose, tribasic calcium phosphate, silicon dioxide, corn starch, sodium starch glycolate, hydroxypropyl cellulose, hydroxypropyl methylcellulose, gelatin, magnesium stearate, microcrystalline cellulose, mannitol or sorbitol, excluding fatty acids and antioxidants. Preferably when a pharmaceutical excipient blend, as defined herein, is utilized in a granulation, said granulation will also contain a substituted 3,5-diene steroidal compound.

In the presently invented methods and in the presently invented tablet formulations preferred antioxidants for use herein are butylated hydroxyanisol (BHA) and butylated hydroxytoluene (BHT), most preferably BHT, preferably in an amount of from about 0.05% to about 2% by weight, more preferably from about 0.1% to about 1% by weight, most preferably about 0.25% by weight.

Contemplated equivalents of substituted 3,5-diene steroidal compounds for use herein are compounds of the formula:



wherein W is an electron-withdrawing group; and the C₁ and C₂ positions and the C₈ to C₁₀ positions are optionally substituted with pharmaceutically acceptable substituents, degrees of unsaturation, a combination of pharmaceutically acceptable substituents and degrees of unsaturation or form part of a larger multicyclic
 5 compound.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative and
 10 not a limitation of the scope of the present invention.

All of the excipients utilized herein are standard pharmaceutical grade excipients available from numerous manufactures well known to those in the art, except that pure stearic acid and pure palmitic acid (from Experiment 2), are not
 15 considered pharmaceutical grade and were obtained from the Aldrich Chemical Company Milwaukee, Wisconsin. Standard grade pharmaceutical excipients, from any reputable source, are considered suitable for use herein.

EXAMPLE 1 - dry granulation process

The lactose, microcrystalline cellulose, sodium starch glycolate, stearic acid, magnesium stearate and N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid shown in Table 10 below, are blended in the proportions shown. The dry
 20 granules are screened and compressed into tablets. Said tablets are optionally coated with, for example, a film coat. The procedure of Example I will produce approximately 750,000 tablets of approximately 133mg each containing about 20
 25 mg of the active ingredient.

TABLE 10

Ingredients	Amounts
N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid	15,000 g
Microcrystalline Cellulose	15,000 g
Lactose	62,500 g
Sodium Starch Glycolate	5,000 g
Stearic Acid	2,000 g
Magnesium Stearate	500 g

EXAMPLE 2 - wet granulation process

The lactose; sodium starch glycolate; microcrystalline cellulose; and N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid shown in Table 11 below, are mixed and granulated in the proportions shown with a 9% solution of the hydroxypropylmethyl cellulose. The wet granules are screened, dried, mixed with the stearic acid; magnesium stearate and butylated hydroxytoluene, screened and compressed into tablets. Said tablets are optionally coated with for example a film coat. The procedure of Example 2 will produce approximately 374,000 tablets of approximately 267mg each containing about 80 mg of the active ingredient.

10

TABLE 11

Ingredients	Amounts
N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid	30,000 g
Lactose	25,000 g
Sodium Starch Glycolate	5,000 g
Microcrystalline Cellulose	32,000 g
Hydroxypropylmethyl Cellulose	4,000 g
Stearic Acid	3,000 g
Magnesium Stearate	500 g
Butylated hydroxytoluene	500 g

While the preferred embodiments of the invention are illustrated by the above, it is to be understood that the invention is not limited to the precise instructions herein disclosed and that the right to all modifications coming within the scope of the following claims is reserved.

15

What is claimed is:

1. A method of preparing stabilized tablet formulations of substituted 3,5-diene steroidal compounds which comprises:
 - 5 (a) admixing a fatty acid, a pharmaceutical excipient blend and a substituted 3,5-diene steroidal compound;
 - (b) compressing the resulting mixture into tablets.
2. The method of claim 1 in which the substituted 3,5-diene steroidal
10 compound is N-t-butyl-androst-3,5-diene-17 β -carboxamide-3-carboxylic acid.
3. The method of claim 1 wherein the fatty acid is stearic acid.
4. The method of claim 3 wherein stearic acid is present in an amount
15 from about 0.5% to about 10% by weight.
5. The method of claim 3 wherein stearic acid is present in an amount from about 1% to about 3% by weight.
- 20 6. The method of claim 3 wherein stearic acid is present in an amount of about 2% by weight.
7. A pharmaceutical formulation in tablet form comprising a pharmaceutical excipient blend, a substituted 3,5-diene steroidal compound and a
25 fatty acid.
8. The pharmaceutical formulation of claim 7 wherein the fatty acid is stearic acid.
- 30 9. The pharmaceutical formulation of claim 8 wherein stearic acid is present in an amount from about 0.5% to about 10% by weight.
10. The pharmaceutical formulation of claim 8 wherein stearic acid is present in an amount from about 1% to about 3% by weight.
35
11. The pharmaceutical formulation of claim 8 wherein stearic acid is present in an amount of about 2% by weight.

12. A method of preparing stabilized tableted formulations of substituted 3,5-diene steroidal compounds which comprises:

- 5 (a) admixing an antioxidant, a pharmaceutical excipient blend, and a substituted 3,5-diene steroidal compound;
(b) compressing the resulting mixture into tablets.

13. The method of claim 12 wherein the antioxidant is butylated hydroxytoluene.

10

14. The method of claim 13 wherein butylated hydroxytoluene is present in an amount from about 0.05% to about 2% by weight.

15 15. The method of claim 13 wherein butylated hydroxytoluene is present in an amount from about 0.1% to about 1% by weight.

16. The method of claim 13 wherein butylated hydroxytoluene is present in an amount of about 0.25% by weight.

20 17. A pharmaceutical formulation in tablet form comprising a pharmaceutical excipient blend, a substituted 3,5-diene steroidal compound and an antioxidant.

25 18. The pharmaceutical formulation of claim 17 wherein the antioxidant is butylated hydroxytoluene.

19. The pharmaceutical formulation of claim 18 wherein butylated hydroxytoluene is present in an amount from about 0.05% to about 2% by weight.

30 20. The pharmaceutical formulation of claim 18 wherein butylated hydroxytoluene is present in an amount from about 0.1% to about 1% by weight.

21. The pharmaceutical formulation of claim 18 wherein butylated hydroxytoluene is present in an amount of about 0.25%.

35

22. A method of preparing stabilized tableted formulations of substituted 3,5-diene steroidal compounds which comprises:

- (a) admixing an antioxidant, a fatty acid, a pharmaceutical excipient blend and a substituted 3,5-diene steroidal compound;
- (b) compressing the resulting mixture into tablets.

5 23. The method of claim 22 wherein the fatty acid is stearic acid and the antioxidant is butylated hydroxytoluene.

 24. The method of claim 23 wherein stearic acid is present in an amount from about 0.5% to about 10% by weight and butylated hydroxytoluene is present
10 in an amount from about 0.05% to about 2% by weight.

 25. The method of claim 23 wherein stearic acid is present in an amount from about 1% to about 3% by weight and butylated hydroxytoluene is present in an amount from about 0.1% to about 1% by weight.
15

 26. The method of claim 23 wherein stearic acid is present in an amount of about 2% by weight and butylated hydroxytoluene is present in an amount of about 0.25%.

20 27. A pharmaceutical formulation in tablet form comprising a pharmaceutical excipient blend, a substituted 3,5-diene steroidal compound, a fatty acid and an antioxidant.

 28. The pharmaceutical formulation of claim 27 wherein the fatty acid is
25 stearic acid and the antioxidant is butylated hydroxytoluene.


 29. The pharmaceutical formulation of claim 28 wherein stearic acid is present in an amount from about 0.5% to about 10% by weight and butylated hydroxytoluene is present in an amount from about 0.05% to about 2% by weight.
30

 30. The pharmaceutical formulation of claim 28 wherein stearic acid is present in an amount from about 1% to about 3% by weight and butylated hydroxytoluene is present in an amount from about 0.1% to about 1% by weight.

35 31. The pharmaceutical formulation of claim 28 wherein stearic acid is present in an amount of about 2% and butylated hydroxytoluene is present in an amount of about 0.25%.

32. The use of stearic acid as a stabilizing agent in tablet formulations of substituted 3,5-diene steroidal compounds.

- 5 33. The use of stearic acid as a stabilizing agent in tablet formulations of compounds of the formula:



- 10 wherein W is an electron-withdrawing group; and the C₁ and C₂ positions and the C₈ to C₁₀ positions are optionally substituted with pharmaceutically acceptable substituents, degrees of unsaturation, a combination of pharmaceutically acceptable substituents and degrees of unsaturation or form part of a larger multicyclic compound.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/09945

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 9/20, 31/565, 47/00

US CL : 514/169,970; 424/465

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/169,970; 424/465

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DERWENT, CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US,A, 4,473,564 (DEWINTER ET AL) 25 SEPTEMBER 1986, COLUMN 10, LINE 60 TO COLUMN 11 LINE 5; COLUMN 11, LINE 17.	12-31
T	WO,A, 93/19758 (HIEBLE ET AL) 14 OCTOBER 1993.	1-11, 32, 33
X	WO,A, 92/00010 (RANDALL) 09 JANUARY 1992, EXAMPLE 2, PAGE 12.	1-11
—		-----
Y		32,33
X	WO,A, 91/13550 (HENRY) 19 SEPTEMBER 1991, EXAMPLE 2, PAGE 19.	1-11
—		-----
Y		32,33
Y	US,A, 4,172,132 (DRAPER ET AL) 23 OCTOBER 1979, COLUMN 14, LINE 37; COLUMN 15, LINE 15-32.	12-31

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

Special categories of cited documents:	
A document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	*F* document member of the same patent family

Date of the actual completion of the international search

19 DECEMBER 1994

Date of mailing of the international search report

DEC 30 1994

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Authorized officer

SHEP ROSE

Facsimile No. (703) 305-3230

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/09945

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US,A, 4,185,101 (DRAPER ET AL) 22 JANUARY 1980, COLUMN 12, LINE 8; COLUMN 12, LINE 50 TO COLUMN 13, LINE 2.	12-31
Y	EP, A, 0,173,478 (HORROBIN) 03 MAY 1986, PAGE 9, LINE 70; PAGE 7, LINES 12-28; PAGE 11, LINE 25.	12-31
Y	US,A, 4,159,326 (BARTON ET AL) 26 JUNE 1979, COLUMN 6, LINES 39-41; COLUMN 10, LINE 54; COLUMN 11, LINES 3-20.	12-31

5739

#5

THIS PAGE BLANK (USPTO)

BEST AVAILABLE COPY
THIS PAGE BLANK (USPTO)